Plant Sociology 59(2) 2022, 27–37 | DOI 10.3897/pls2022592/03 *Topical Collection: "Conservation and biodiversity of bryophytes"*

PENSOFT

Società Italiana di Scienza della Vegetazione (SISV)

The bryophyte vegetation of the gypsum outcrops of Sicily

Marta Puglisi[1], Maria Privitera[1], Giovanni Spampinato[2]

1 *Department of Biological, Geological and Environmental Sciences, University of Catania, Italy*
2 *Department of Agricultural Science, Mediterranean University of Reggio Calabria, Italy*

Corresponding author: Marta Puglisi (mpuglisi@unict.it)

Subject editor: Silvia Poponessi ♦ **Received** 29 July 2022 ♦ **Accepted** 4 October 2022 ♦ **Published** 7 December 2022

Abstract

A phytosociological study on the gypsicolous bryophyte communities of Sicily was carried out. The surveyed communities are: *Tortuletum revolventis*, *Trichostomo crispuli-Tortuletum revolventis*, *Crossidio squamiferi-Aloinetum aloidis* of the order *Barbuletalia unguiculatae*, and *Crossidio crassinervis-Tortuletum obtusatae* of the order *Tortulo brevissimae-Aloinetalia bifrontis*. The associations were examined from a synecological, synhierarchical and chorological point of view. A life form and life strategy analysis of all communities reflect the response of plant functional types towards the environmental demands. Only one life strategy dominates the communities; acrocarpous, turf-forming colonists clearly prevail on gypsum outcrops, subject to drought stress. They provide the main functional type within pioneering communities and communities of first successional stages. The keystone characters outlined ensure a successful dispersal, establishment and habitat maintenance of the species and associations.

Keywords

bryophytes, gypsum, life forms, life strategies, phytosociology, Sicilian territory

Introduction

Gypsum crops out in almost all Italian regions, with the main location on the northern border of the Apennine chain of the Emilia Romagna region and in Sicily; the last one comprises the largest gypsum outcrops with a total area of more than 1,000 km^2 followed by Emilia Romagna with a surface more than 100 km^2 (Forti and Sauro 1996). Most Italian gypsum outcrops belong to the Gessoso-Solfifera Formation (GSF) formed during the "Messinian salinity crisis" (MSC), caused by reduced water inflow from the Atlantic Ocean to the Mediterranean Sea with the consequent salt precipitation and decrease in Mediterranean level due to evaporation. Sicily, due to its geographical location in the centre of the Mediterranean, is the region where the Gessoso-Solfifera Formation crops out more extensively with the highest development in the provinces of Caltanissetta, Enna and Agrigento.

In gypsum substrates there is an incipient impoverishment of the soils, especially in arid and semi-arid areas where the high evaporation leads to a capillary rise of the gypsum towards the surface layers, inducing recrystallization of the gypsum and the formation of crusts (Escudero et al. 2015). The gypsum crusts are inhospitable substrates to higher plants; only cyanobacteria, lichens and, with difficulty, few bryophytes can survive (Escudero et al. 2015). Gypsum has low water retention to which a high osmotic pressure of the soil solution is added which strongly reduces root absorption in higher plants (Herrero and Porta 2000), not in bryophytes which are, instead, without absorbent roots. For all these characteristics related to the type of substrate and for the environmental dryness, the plant component is poor, selected, and specialized.

Gypsum outcrops are peculiar habitats of naturalistic interest and, despite the inhospitality of the substrate, host a valuable plant diversity with many specialized and rare species, as well as some interesting communities according to the peculiar characteristics of substrate and climatic conditions.

As regards the bryoflora, the knowledge in Italy is quite good, even if not exhaustive; the contributions regard the gypsum outcrops of six sites of the Nature 2000 Network of the Emilia Romagna Region (Aleffi et al. 2014) and the main gypsum areas of Sicily (Puglisi et al. 2019); in the

Copyright *Marta Puglisi et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

last paper the functional group of gypsophily was also attributed to all surveyed species.

As concerns the bryophyte vegetation, the knowledge is limited to some gypsum areas of central Sicily (Privitera 1989); other sporadic data can be found in Lo Giudice and Bonanno (2010). Therefore, this limited knowledge, together with the interest generated by the peculiarity of the habitat, has led us to undertake an extensive investigation on the bryophyte communities colonizing the gypsum outcrops.

Methods

Study areas

The most complete succession of the Gessoso-Solfifera Formation crops out in Sicily. Here, the temporary closure of the Mediterranean Basin led to the creation of two different deposition cycles, one upper and one lower. In fact, two evaporitic successions can be recognized, laying upon pelitic facies of the Terravecchia Formation and locally on white diatomitic marls (Tripoli Formation), and covered by fine pelagic sediments (Trubi Formation), (De Waele et al. 2017). The lower evaporitic unit consists of evaporitic limestone, crystalline gypsum interlayered with gypsy marls and salts (mainly chlorides). The upper evaporitic unit consists of crystalline gypsum, primary or clastic, bioclastic limestone and, finally, clayey sands (Catalano 1986). The gypsum is poorly permeable substrate, very friable and subject to erosive processes that give a "young" character, preventing the formation of deep and advanced soils.

The Sicilian territory is characterized by a Mediterranean climate, diversified based on the altitudes and slopes. The average annual temperature is 17-18° C in the coastal areas, and 16° C in the most inland areas. Precipitation is concentrated in autumn and winter with an average of 500–700 mm/year. According to the bioclimatic classification of Rivas-Martínez et al. (2011), the bioclimate of the investigated sites is of the Mediterranean pluviseasonal oceanic, ranging from lower thermomediterranean to lower mesomediterranean thermotype and from lower dry to upper semiarid ombrotype (Pesaresi et al. 2014).

The phanerogamic vegetation structured by gypsophytes grow directly on gypsum outcrops or on thin layers of lithosol, poor-nutrient in low productive environments, not on the soils deriving from the gypsum where it is possible to find the scrub of the *Oleo sylvestris-Ceratonion siliquae* Br.-Bl. ex Guinochet & Drouineau 1944, probably due to predominant type bioclimate (Mediterranean sub-humid) that cause a prevalent flow-down movement of water in the soil profile and give rise to relatively fertile soils. The most important plant communities on vertical or sub-vertical gypsum rocks are *Brassica villosa* (Biv.) subsp. *tineoi* (Lojac.)Raimondo & Mazzola communities and *Diplotaxis harra* (Forssk.)Boiss subsp. *crassifolia* (Raf.)Maire communities, referable to *Dianthion rupicolae* Brullo & Marcenò 1979, and ephemeral vegetation of *Sedo-Ctenopsion gypsophilae* Rivas Goday & Rivas-Martínez ex Izco 1974 on slightly steep gypsum outcrops with three different communities: *Filagini-Chaenorhinetum rupestris* Brullo, Marcenò, Minissale & Spampinato 1989, *Sedum gypsicola* subsp. *trinacriae* community and *Petrosedum ochroleucum* subsp. *mediterraneum* community.

Sampling methods

The phytosociological study, which follows the plant sociological method of Braun-Blanquet (1964), is based on unpublished and literature data. As regards the unpublished data, the field work was carried out during 2019-2021 years; the season of the collection was spring or late spring. The cover of each taxon is estimated according to the following values: + (<1%), 1 (1-10%), 2 (10.1-25%), 3 (25.1-50%), 4 (50.1-75%), 5 (75.1-100%).

The surveys and samplings were done on the gypsum outcrops in sites referred to the provinces of Caltanissetta, Palermo, Trapani and Agrigento. The literature data refer to sites of the Enna and Catania provinces and, partially, to Caltanissetta and Agrigento, too (Privitera 1989; Lo Giudice and Bonanno 2010; Puglisi et al. 2022). The list of the sites (Fig. 1) is reported below.

1. Rocca di Entella (municipality of Contessa Entellina), gypsum crusts, bioclimate upper thermomediterranean upper dry.
2. Castelluccio (municipality of Contessa Entellina), selenite gypsum, bioclimate upper thermomediterranean upper dry.
3. Serre di Ciminna (municipality of Ciminnà), selenite gypsum, bioclimate lower mesomediterranean upper dry.
4. S. Ninfa (municipality of S. Ninfa), selenite gypsum, bioclimate upper thermomediterranean upper dry.
5. Sutera outskirts, selenite gypsum, bioclimate upper thermomediterranean lower dry.
6. Milena hills, selenite gypsum, bioclimate upper thermomediterranean lower dry.
7. Enna outskirts (from Lo Giudice and Bonanno 2010), gypsum outcrops, bioclimate upper mesomediterranean lower subhumid.
8. Raddusa outskirts (from Lo Giudice and Bonanno 2010), gypsum outcrops, bioclimate lower mesomediterranean lower dry.

Some of the investigated sites fall within within protected areas, such as Integral Nature Reserve "Grotta di Entella" (site 1 and site 2), Oriented Nature Reserve "Serre di Ciminna" (site 3), Integral Nature Reserve "Santa Ninfa" (site 7), and/or within the SAC of Natura 2000 network: ITA020024 Rocche di Ciminna (site 3), ITA020042 Rocche di Entella (site 1 and site 2), ITA010022 Complesso Monti di Santa Ninfa – Gibellina e Grotta di Santa Ninfa (site 7).

The identification of the outcrops was carried out through the preliminary analysis of the geological map

of Sicily (Lentini and Carbone 2014); moreover, the field delimitation of the gypsum outcrops was carried out considering the vascular gypsophilous vegetation as practical guide for recognition of gypsum soil.

For the synecological analysis, life forms and life strategies of each taxon were considered. The life forms follow the concept of Mägdefrau (1982), those of the life strategies During (1979), Frey and Kürschner (1991), Kürschner and Frey (2013). For each species and category the mean percentage cover (MPC) values of the biological parameters within the association is calculated, based on the cover values (Puglisi et al. 2013, 2014, 2018).

The syntaxonomic arrangement and nomenclature follow Puglisi and Privitera (2012); the nomenclature of the taxa follows Hodgetts et al. (2020).

Results and discussion

Based on the phytosociological relevés, the investigated bryophyte vegetation is represented by xerophytic, basiphytic communities, found mostly on selenite outcrops in small depressions or concavities of rocks with accumulated soil. These communities are largely dominated by acrocarpous, small-sized mosses with a turf biotype and xero-pottioid life syndrome; conversely, the pleurocarpous mosses and liverworts are very rare. In more xeric areas, only some lichens, particularly resistant to the harshness of the chalky substrate, can be found.

From a phytosociological point of view, these pioneer bryophyte communities are referred to the class *Psoretea decipientis*. The surveyed communities are discussed below.

TORTULETUM REVOLVENTIS Marstaller 1980 (Tab. 1)

It is the most widespread and typical association characterizing the gypsum outcrops of Sicily. Its presence was highlighted in many localities with a thermomediterranean and mesomediterranean dry rarely subhumid bioclimates, such as Serre di Ciminna, Castelluccio, S. Ninfa, Sutera outskirts, Milena hills, Raddusa outskirts, and in some gypsum outcrops of Enna. The association can be defined as xerophytic, basiphytic, photophytic and strictly gypsophytic. It was found in small depressions where a thin layer of soil accumulates, as well as in the cracks between the gypsum crystals in the ambit of the association. The cover ranges from 30% to 100%; the surfaces are 20-50 dm^2. The association is floristically characterized by *Tortula revolvens* (Schimp.)G. Roth, occurring with high cover values only in few relevès, in the sites of Enna and Catania, in more mesic (less edaphic drought) conditions. This species is associated with some characteristics of the alliance *Grimaldion fragrantis* and a lot of species of the order *Barbuletalia unguiculatae* and class *Psoretea decipientis* with the most represented *Didymodon vinealis* and *Barbula unguiculata*. In the sites of Serre di Ciminna and S. Ninfa, in more edaphic drought conditions, it is possible to find some transgressive species of the order *Tortulo brevissimae-Aloinetalia bifrontis*, and a lichen component



Figure 1. Location of the investigated sites.

Table 1. *Tortuletum revolventis* Marstaller 1980.

Relevé number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31
Altitude (m a.s.l.)	510	505	505	410	410	400	400	415	320	380	300	400	400	350	350	400	320	320	300	750	750	750	750	740	740	330	330	330	330	480	480
Surface (dm²)	20	30	20	20	20	20	30	15	40	50	30	30	20	40	40	20	30	20	30	12	25	20	30	20	15	25	20	15	22	30	40
Cover (%)	50	35	65	65	45	40	30	65	30	40	65	60	55	75	70	30	50	65	70	100	70	50	50	100	70	70	40	100	60	35	40
Number of species	8	9	9	11	9	8	6	6	5	5	5	4	4	5	6	3	4	5	7	9	7	9	9	14	9	7	7	7	8	8	7
Char. association																															
Tortula revolvens (Schimp.) G.Roth	2	1	2	3	2	2	1	3	2	2	3	3	3	4	3	2	3	3	2	4	3	2	2	4	3	3	2	3	2	1	1
Char. alliance, order and class (*Grimaldion fragrantis*, *Barbuletalia unguiculatae*, *Psoretea decipientis*)																															
Barbula unguiculata Hedw.	1*	1	.	2	.	1	.	+	+	.	.	.	.	.	.	1	1	2	1	.	.	1	+	+	1	.	.	.	+	.	2
Didymon vinealis (Brid.) R.H.Zander	.	+	2	1	.	1	1	.	.	.	.	+	.	.	.	.	.	1	1	.	.	.	+	2	.	.	.	1	1	.	1
Didymon luridus Hornsch.	.	.	.	+	+	.	+	1	.	+	+	.	.	.	.	.	.	.	.	+	+	1	.	+	1	1	.	.	.	.	.
Encalypta vulgaris Hedw.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	2	2	1	2	1	1	2	1	2	1	.	.
Pseudocrossidium hornschuchianum (Schultz) R.H.Zander	.	1	+	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	+	1	+	+	+	+	.	.	.	.
Didymodon acutus (Brid.) K.Saito	+	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	+	.	.	+	1	.	2	+
Weissia controversa Hedw. var. *controversa*	.	.	.	+	+	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	1	+	.	+	.	.	1	.
Ptychostomum capillare (Hedw.) Holyoak & N.Pedersen	2	.	1	1	1	+	.	2	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Bryum radiculosum Brid.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	+	.	.	+	+	.	.	.	.	1
Bryum dichotomum Hedw.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	+	.	.	+	1
Aloina ambigua (Bruch & Schimp.) Limpr.	.	.	.	.	1	+	.	.	.	+	.	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Microbryum starkeanum (Hedw.) R.H.Zander	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	+	.	.	.	.	+	.	.
Trichostomum brachydontium Bruch	.	.	1	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Trichostomum crispulum Bruch	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	.	1	.	.	.
Fissidens viridulus (Sw.)Wahlenb.	+	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Streblotrichum convolutum (Hedw.) P.Beauv. var. *convolutum*	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1
Trasngressive of the order *Tortulo brevissimae-Aloinetalia bifrontis*																															
Fossombronia caespitiformis (Raddi) De Not. ex Rabenh. subsp. *caespitiformis*	.	1	1	.	.	+	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Dicranella howei Renauld & Cardot	+	.	.	+	1	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Crossidium crassinervium (De Not.)Jur.	.	.	.	.	.	.	.	.	1	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Microbryum davallianum (Sm.) R.H.Zander var. *davallianum*	.	.	.	+	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Other species																															
Pseudocrossidium revolutum (Brid.) R.H.Zander	.	.	1	1	.	.	1	.	+	+	2	.	.	1	+	2	.	.	.	.	+	.	.	.	.	.	.	.	.	1	.
Ptychostomum imbricatulum (Müll.Hal.) Holyoak & N.Pedersen	.	1	2	.	.	.	.	.	.	+	.	.	.	.	2	.	.	.	.	1	.	.	.	+	.	.	.	+	+	.	.
Gymnostomum viridulum Brid.	.	.	.	.	.	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	1	+	2	3	2	.	.
Didymodon rigidulus Hedw.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	+	+	.	.	+	+	.	.	+	.	.
Crossidium squamiferum (Viv.)Jur. var. *squamiferum*	.	.	.	.	.	.	.	.	.	.	.	1	1	.	.	.	.	.	.	+	.	+	.	.	.	.	.	.	.	.	.
Ptychostomum donianum (Grev.) Holyoak & N.Pedersen	.	.	.	.	.	.	.	.	+	.	1	.	.	1	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Didymodon insulanus (De Not.) M.O.Hill	1	.	+	.	.	.	+	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.
Brachytheciastrum velutinum (Hedw.) Ignatov & Huttunen	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	.	+	+	+	.	.	.	.	.	.	.	.	.	.	.	.
Tortella squarrosa (Brid.)Limpr.	.	.	.	.	.	.	.	.	.	.	.	.	+	.	.	.	.	.	.	.	.	.	+	+	.	.	.	.	.	.	.

represented by *Lathagrium cristatum* (L.) Otálora, P.M. Jørg. & Wedin and *Gyalolechia subbracteata* (Nyl.) Søchting, Frödén & Arup. Moreover, the relevés from Enna and Raddusa outskirts, characterized by the occurrence of *Encalypta vulgaris* Hedw., could represent a more mesophytic variant of the association.

Tortuletum revolventis, described for the chalky substrates of Thuringia (central Germany) (Marstaller 1980, 2008), is already known for some of the gypsum areas of central Sicily (Privitera 1989).

TRICHOSTOMO CRISPULI-TORTULETUM REVOLVENTIS Puglisi, Spampinato & Privitera 2022 (Tab. 2)

The community was found on selenite gypsum in north-western and central Sicily at Castelluccio, Sutera outskirts, Milena hills. It is a gypsycolous, terricolous, photo-sciophytic, meso-xerophilous, covering small surfaces such as concavities of rocks with a conspicuous layer of soil, making it the least pioneer among the structured communities with *Tortula revolvens*. *Trichostomo crispuli-Tortuletum revolventis* was found in the thermomediterranean dry bioclimatic belt, in more protected and less xeric conditions than *Tortuletum revolventis*. The surface cover ranges from 5 to 20 dm^2 and the sites are mostly flat or scarcely inclined; the cover varies between 55% and 95%.

The community is characterized by *Trichostomum crispulum* Bruch, having major cover values, and *Tortula revolvens*, constantly present. To these species a lot of characteristics of higher units (*Grimaldion fragrantis*, *Barbuletalia unguiculatae*, *Psoretea decipientis*) are associated too. This community differs from *Tortuletum revolventis* in colonizing more protected and less exposed sites, and for its less pioneering character. *Tortula revolvens* (incl. var. *obtusata*) is also indicated as a characteristic of *Crossidio crassinervis-Tortuletum obtusatae* Ros & Guerra 1987 and *Trichostomopso-Tortuletum obtusatae* Frey, Herrnstadt & Kürschner 1990 of the order *Tortulo brevissimae-Aloinetalia bifrontis* but differs from both for the less xeric character and the different floristic set. The community shows some affinity with the association *Trichostomo crispuli–Aloinetum* aloides from which it differs for the overall floristic composition and for the gypsophytic character.

CROSSIDIO SQUAMIFERI-ALOINETUM ALOIDIS Guerra & Varo 1981 (Tab. 3)

This association was found in the thermomediterranean and mesomediterranean bioclimatic belts at Sutera outskirts, Milena hills and Raddusa outskirts; in these sites, it colonized gypsum rocky fissures with accumulated soil. The association behaves as chasmochomophytic, basiphyticous, xerophytic, photo-sciophytic. This association is not strictly linked to gypsum substrates, being found also on marl, marly-gypsum and clayey soil, limestone rocky fissures (Privitera and Puglisi 1999; Lo Giudice and Bonanno 2010), behaving as not strictly gypsophytic. The surfaces are quite small (4–6 dm^2) and the cover varies between 30% and 65%. Floristically, it is characterized by *Crossidium squamiferum* (Viv.)Jur. var. *squamiferum*, a Circum-Tethyan species, and *Aloina aloides* (Koch ex Schultz) Kindb., accompanied by a set of characteristics of the alliance *Grimaldion fragrantis*, order *Barbuletalia unguiculatae* and class *Psoretea decipientis*. *Crossidio squamiferi-Aloinetum aloidis* is known in Italy only in Sicily and southern Calabria (Privitera and Puglisi 1999; Lo Giudice and Bonanno 2010).

Table 2. *Trichostomo crispuli-Tortuletum revolventis* Puglisi, Spampinato & Privitera 2022.

Relevé number	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Altitude (m a.s.l.)	477	477	464	464	452	452	305	305	315	345	345	210	210	195
Cover (%)	80	65	90	80	60	70	95	90	55	85	85	85	70	95
Surface (dm^2)	7	20	10	8	8	10	5	10	20	10	8	10	20	8
Inclination (°)	15	-	-	30	-	-	20	-	-	-	-	-	20	-
Exposure	SW	-	-	SE	-	-	S	-	-	-	-	-	S	-
Number of species	4	8	5	4	5	5	5	6	6	6	6	5	4	5
Char. association														
Trichostomum crispulum	4	2	4	4	3	3	5	4	3	4	4	4	3	4
Tortula revolvens	2	1	3	2	1	1	2	2	1	2	2	2	2	3
Char. alliance, order and class (*Grimaldion fragrantis, Barbuletalia unguiculatae, Psoretea decipientis*)														
Trichostomum brachydontium	1	1	.	+	1	.	.	1	.	1	1	1	.	+
Didymon vinealis	.	.	.	.	.	.	1	2	1	.	+	.	2	1
Didymodon acutus	.	1	+	1	.	2	.	.	+	.	.	.	.	.
Weissia controversa var. *controversa*	.	.	.	.	.	.	1	.	+	.	1	1	.	.
Bryum dichotomum	.	1	1	.	.	1	.	.	.	.	.	.	.	.
Streblotrichum convolutum var. *convolutum*	.	2	.	.	.	1	.	.	.	.	.	.	.	.
Aloina ambigua	.	.	.	.	.	.	.	1	.	+	.	.	.	.
Didymon luridus	.	.	.	.	.	.	.	.	.	1	+	.	.	.
Other species														
Homalothecium sericeum (Hedw.) Schimp.	+	.	+	.	1	.	.	+	1	+	.	+	+	1
Tortula solmsii (Schimp.)Limpr.	.	1	.	.	.	.	.	.	.	.	.	.	.	.
Tortella inflexa (Bruch)Broth.	.	1	.	.	.	.	.	.	.	.	.	.	.	.
Petalophyllum ralfsii (Wilson) Nees & Gottsche	.	.	.	.	1	.	.	.	.	.	.	.	.	.
Tortella squarrosa	.	.	.	.	.	.	+	.	.	.	.	.	.	.

CROSSIDIO CRASSINERVIS-TORTULETUM OBTUSATAE Ros & Guerra 1987 (Tab. 4)

This community grows on very dry, very exposed, soil with a thin layer of gypsum at Rocca di Entella in northern Sicily in the thermomediterranean dry bioclimatic belt. From an ecological point of view, it can be considered as a terricolous, thermophytic, markedly xerophytic, photophytic community. It finds its optimum in arid territories of the Mediterranean and Irano-Turanian region (García-Zamora et al. 2000). The cover is not high, ranging from 30% to 65 %; the surfaces are 10 to 30 dm^2. Characteristic species of the association are *Tortula revolvens* and *Crossidium crassinervium*, both species with main distribution areas in the Mediterranean, Irano-Turanian and Saharo-Arabian territories. As concerns *Tortula revolvens*, in the description of the association (Ros and Guerra 1987), the characteristic species is *Tortula revolvens* var. *obtusata* Reimers, considered now a synonym of *Tortula revolvens* (Schimp.) G. Roth. To these species a set of characteristics of the alliance *Aloino bifrontis-Crossidion crassinervis* and order *Tortulo brevissimae-Aloinetalia bifrontis* are associated, such as *Dicranella howei* Renauld & Cardot, *Fossombronia caespitiformis* (Raddi) De Not. ex Rabenh. subsp. *caespitiformis*, *Didymodon tophaceus* (Brid.)Lisa subsp. *sicculus* (M.J.Cano, Ros, García-Zam. & J.Guerra) Jan Kučera, *Microbryum davallianum* (Sm.) R.H.Zander var. *davallianum*, *Tortula brevissima* Schiffn. and *Tortula atrovirens* (Sm.)Lindb. Among the other species, a group of lichens are found too, such as *Lathagrium cristatum*, *Squamarina lentigera* (Weber)Poelt and *Gyalolechia subbracteata*. From a phytosociological point of view, *Crossidio crassinervis-Tortuletum obtusatae* is included in the above-mentioned syntaxa, which group markedly xerophytic, photophytic communities with Mediterranean, Irano-Turanian and Saharo-Arabian distribution The association is signalled from Spain (Ros and Guerra 1987; Guerra et al. 1993; Moya et al. 1994; García-Zamora et al. 2000) and Israel (Frey et al. 1990), where it was found on gypsum and marl-gypsum substrates. *Crossidio crassinervis-Tortuletum obtusatae* is reported for the first time for the bryophyte vegetation of Italy.

Plant functional types

The plant functional types (life forms, life syndromes, morphological-anatomical keystone characters) characterizing the communities of the Sicilian gypsum substrates provide useful information on the mechanisms of habitat maintenance, establishment, re-establishment and dispersal of the species and communities (e.g., Kürschner and Frey 2013, Puglisi et al 2014, 2015, 2016a, 2016b, 201). The main characters used for the analysis (Tab. 5) are life span (avoidance vs. tolerance strategy of the gametophyte), breeding system, main reproductive effort (sexual vs. asexual reproduction) and dispersal strategies [small spores (< 25μm) providing chance dispersal vs. large spores (> 25 μm) indicating decreasing long-range dispersal and achory].

The communities show a relatively low diversity concerning plant functional types (life forms and life strategies), with a very high prevalence of the colonist strategy with MPC values of 90.7% in *Tortuletum revolventis*, 91.0% in *Crossidio crassinervis-Tortuletum obtusatae*, 94.7% in *Trichostomo crispuli-Tortuletum revolventis* and 95.7% in *Crossidio squamiferi-Aloinetum aloidis* (Tab. 6). Many investigations have shown that communities of harsh habitats, especially those of sunny sites, rocks, as

Table 3. *Crossidio squamiferi-Aloinetum aloidis* Guerra & Varo 1981.

Relevés number	1	2	3	4	5	6	7
Altitude (m a.s.l.)	375	380	365	380	385		
Cover (%)	40	65	40	50	30	30	40
Surface (dm2)	4	6	5	5	5	6	6
Inclination (°)	40	30	60	40	40	30	30
Exposure	SW	SE	E	NW	SW	SW	E
Species number	6	4	5	5	3	4	6
Char. association							
Aloina aloides (Koch ex Schultz) Kindb.	1	.	+	1	.	.	1
Crossidium squamiferum var. *squamiferum*	1	3	2	2	2	2	2
Char. alliance, order and class (*Grimaldion fragrantis, Barbuletalia unguiculatae, Psoretea decipientis*)							
Didymodon vinealis	2	+	.	1	+	1	.
Barbula unguiculata	1	.	.	1	.	+	.
Didymodon luridus	1	.	1	.	.	.	.
Weissia controversa var. *controversa*	.	1	+	.	.	.	.
Aloina ambigua	.	.	.	.	.	.	2
Pseudocrossidium hornschuchianum	.	.	1	.	.	.	.
Ptychostomum capillare	1	.	.	.	.	.	.
Trichostomum brachydontium	.	.	.	.	1	.	.
Other species							
Pseudocrossidium revolutum	.	1	.	1	.	.	.
Ptychostomum torquescens (Bruch & Schimp.) Ros & Mazimpaka	.	.	.	.	.	1	.
Encalypta vulgaris Hedw.	.	.	.	.	.	.	+
Tortula muralis Hedw. supsp. *muralis*	.	.	.	.	.	.	+
Ptychostomum imbricatulum	.	.	.	.	.	.	+

Table 4. Crossidio crassinervis-Tortuletum obtusatae Ros & Guerra 1987.

Relevé number	1	2	3	4	5	6	7	8
Altitude (m a.s.l.)	520	520	525	515	515	505	505	510
Surface (dm²)	20	10	30	15	20	30	20	10
Cover (%)	55	65	45	60	40	50	30	45
Species number	7	7	7	8	7	8	5	4
Char. association								
Tortula revolvens	3	3	2	3	2	2	2	3
Crossidium crassinervium	1	1	2	+	1	1	1	1
Char. alliance and order (*Aloino bifrontis-Crossidion crassinervis, Tortulo brevissimae-Aloinetalia bifrontis*)								
Tortula brevissima Schiffn.	1	.	+	1	1	.	+	1
Dicranella howei	.	+	.	1	.	1	1	.
Fossombronia caespitiformis subsp. *caespitiformis*	1	2	.	+	.	1	.	+
Didymodon tophaceus (Brid.)Lisa subsp. *sicculus* (M.J.Cano, Ros, García-Zam. & J.Guerra) Jan Kučera	+	.	1	.	1	+	.	.
Microbryum davallianum var. *davallianum*	+	1	.	+	.	.	.	.
Tortula atrovirens (Sm.) Lindb.	.	.	.	1	.	1	.	.
Char. class (*Psoretea decipientis*)								
Bryum dichotomum	.	.	+	1	1	.	.	1
Pseudocrossidium hornschuchianum	1	.	.	.	.	.	.	.
Other species								
Pseudocrossidium revolutum	.	+	.	1	1	.	1	.
Aloina brevirostris (Hook & Grev.) Kindb.	.	1	+	.	+	.	.	.
Tortella flavovirens (Bruch) Broth. var. *flavovirens*	.	.	.	1	.	2	.	.
Crossidium squamiferum var. *squamiferum*	1	.	.	.	.	.	+	.
Aloina aloides	.	.	.	.	.	1	.	.
Homalothecium sericeum	.	.	1	.	.	.	.	.
Lathagrium cristatum (L.) Otálora, P.M. Jørg. & Wedin	1	+	.	1	.	.	+	.
Squamarina lentigera (Weber)Poelt	.	.	1	.	.	.	.	.
Gyalolechia subbracteata (Nyl.) Søchting, Frödén & Arup	.	.	.	.	1	.	.	.

Table 5. Characters, life forms and life strategies of the communities. Abbreviations Life forms: Sc: solitary creeping; St: solitary thalloid; Tf turf; Tp: turf protonemal; Ts turf scattered; We weft. Life cycle: a annual/ephemeral; p pauciennial/pluriennial; pe perennial. Sexual reproduction: 1 frequent within the 1st year, 2 frequent within the 2nd – 4th year; [ ] data uncertain; a abundant; a/r absent or rare; f frequent; o occasional; p present. Breeding system: D dioicous; M monoicous. Asexual reproduction: a/r absent or rare; bl brood leaves; fl fragmentation of leaves/caducous leaves; Ge gemmae; Rhg rhizoid gemmae. Dispersal strategy: lr long range dispersal, sr short range dispersal. Life strategies: AnS anniual shuttle; C colonists; PeS perennial shuttle species; PS perennial stayers; PaS short-lived shuttle species; as with high asexual reproductive effort; m with moderate or rare sexual and asexual reproductive effort; s with high sexual reproductive effort.

	Life foms	Life cycle	Sexual reproduction	Breeding system	Spores (Ø in µm)	Asexual reproduction	Dispersal strategy	Life strategies
Aloina aloides	Ts	p	f	D	15-25	a/r	sr,lr	**Cs**
Aloina ambigua	Ts	p	f	D	10-15	a/r	sr,lr	**Cs**
Aloina brevirostris	Ts	p	a/r	D	10-15	a/r	sr,lr	**Cs**
Aloina rigida	Ts	p	a/r	D	14-16	a/r	sr,lr	**Cs**
Streblotrichum convolutum var. *convolutum*	Tf	p	[2]	D	8-10	Rhg	sr,lr	**Cas**
Barbula unguiculata	Tf	p	f	D	10-14	a/r	sr,lr	**Cs**
Brachytheciastrum velutinum	We	pe	2	M	13-16	a/r	sr,lr	**PSas**
Bryum argenteum	Tf	p	1	D	8-14	g	sr,lr	**Cas,s**
Bryum dichotomum	Tf	p	a/r	D	8-16	Ge,Rhg	sr,lr	**Cas**
Bryum radiculosum	Tf	p	f	D	10-14	Rhg	sr,lr	**Cas**
Cephaloziella baumgartneri	Ms	p	f	M	8-12	Ge	sr,lr	**Cas**
Ceratodon purpureus	Tf	p	2	D	10-14	a/r	sr,lr	**Cs**
Crossidium crassinervium	Ts	p	a/r	M	9-16	a/r	sr,lr	**Cm**
Crossidium squamiferum var. *squamiferum*	Tf	p	2	M	9-22	a/r	sr,lr	**Cm**
Dicranella howei	Tf	p	a/r	D	-	Rhg	sr,lr	**Cm**
Didymodon acutus	Tf	p	o	D	9-12	Ge	sr,lr	**Cas**
Didymodon insulanus	Tf	p	a/r	D	~ 10	a/r	sr,lr	**Cm**
Didymodon rigidulus	Tuft	p	a/r	D	10-12	Ge	sr,lr	**Cas**
Didymodon sicculus	Tf	p	a/r	D	-	a/r	sr,lr	**Cm**
Didymon luridus	Tf	p	2	D	12-16	a/r	sr,lr	**Cs**
Didymon vinealis	Tuft	pe	2	D	10-13	a/r	sr,lr	**Cs**
Encalypta vulgaris	Tuft	pe	2	M	30-45	a/r	sr,lr–ac	**PeSs**
Entosthodon convexus	Tuft	a	2	-	18-26	a/r	sr,lr	**F**
Fissidens viridulus	Tf	p,pe	2	M,D	8-15	a/r	sr,lr	**Cs**
Fossombronia caespitiformis subsp. *caespitiformis*	Sc	a,p	1	M	38-56	a/r	sr,lr–ac	**AnS**
Grimmia pulvinata	Cu	p	[1]	M	8-12	a/r	sr,lr	**Cpa**
Gymnostomum viridulum	Tf	p	a/r	D	10-12	a/r	sr,lr	**Cas**
Homalothecium aureum	We	pe	o	D	14-18	a/r	sr,lr	**PSm**
Homalothecium sericeum	Mr	pe	a/r	D	11-22	a/r	sr,lr	**PSm**

Table 5. Continuation.

	Life foms	Life cycle	Sexual reproduction	Breeding system	Spores (Ø in µm)	Asexual reproduction	Dispersal strategy	Life strategies
Microbryum davallianum var. *davallianum*	Ts	a	a	M	31-40	a/r	sr,lr–ac	**Ans**
Microbryum starkeanum	Ts	a	a	M	19-42	a/r	sr,lr–ac	**Ans**
Petalophyllum ralfsii	St	p	f	D	40-56	a/r	sr,lr–ac	**PaS**
Pseudocrossidium hornschuchianum	Tf	p	a/r	D	8-10	a/r	sr,lr	**Cm**
Pseudocrossidium revolutum	Tf	p	a/r	D	10-14	a/r	sr,lr	**Cm**
Ptychostomum capillare	Tf	p	1	D	9-15	Rhg	sr,lr	**Cs,as**
Ptychostomum donianum	Tf	p	a/r	D	12-14	a/r	sr,lr	**Cs**
Ptychostomum imbricatulum	Tf	p	2	D	10-14	Rhg	sr,lr	**Cs,as**
Ptychostomum torquescens	Tf	p	1	D	10-16	Ge,Rhg	sr,lr	**PSs,as**
Scleropodium touretii	Mr	pe	a/r	D	11-18	a/r	sr,lr	**PSm**
Timmiella barbuloides	Tp	pe	2	M	10-13	a/r	sr,lr	**PSs**
Tortella flavovirens var. *flavovirens*	Tf	p	a/r	D	12-14	a/r	sr,lr	**Cas**
Tortella inflexa	Tp	p	a/r	D	8-10	a/r	sr,lr	**Cm**
Tortella squarrosa	Tf	p,pe	a/r	D	10-13	bl	sr,lr	**PSas**
Tortula atrovirens	Tf	p	2	M	24-28	a/r	sr,lr	**Cm**
Tortula brevissima	Ts	p	a/r	D	14-15	a/r	sr,lr	**Cs**
Tortula muralis susp. *muralis*	Tf	pe	1	M	7-14	a/r	sr,lr	**Cs**
Tortula revolvens	Tf	p	a/r	-	8-10	a/r	sr,lr	**Cm**
Tortula solmsii	Tf	pe	o	D	10-15	a/r	sr,lr	**Cm**
Trichostomum brachydontium	Tf	p,pe	a/r	D	14-18	± fl	sr,lr	**PeSm**
Trichostomum crispulum	Tf	p,pe	a/r	D	16-18	a/r	sr,lr	**Cm**
Weissia condensa	Tf	p	o	M	14-24	a/r	sr,lr	**Cm**
Weissia controversa	Tf	p	2	M	16-20	a/r	sr,lr	**Cs**

well as communities growing on anthropogenic sites and as pioneer communities (primary succession stages) are the domain of the colonists life strategy (Kürschner and Frey 2013). Colonists are characterized by a generally low gametophyte longevity (often pauciennial), an often high asexual reproduction by rhizoid gemmae and leaf gemmae for a rapid establishment, a regular formation of sporophytes and the production of numerous small spores (< 25µm in diameter). However, due to the inhospitality of the substrate, the colonist species of the gypsophytic communities show a moderate reproductive effort, with MPC values ranging from 58.4% in to 83.4% (Tab. 6).

Of no significant value are the shuttle and the perennial stayers strategies, with very low MPC incidence (Tab. 6).

Strongly correlated to the colonist strategy are the life forms turf and tuft, as typical for many acrocarpous mosses, with MPC values of the turf (including the turf scattered and turf protonemal) corresponding to 94.0% in *Tortuletum revolventis*, 93.9% in *Trichostomo crispuli-Tortuletum revolventis*, 88.5% in *Crossidio squamiferi-Aloinetum aloidis* and 92.3 in *Crossidio crassinervis-Tortuletum obtusatae*. The percentage of the other life forms is negligible.

Conclusion

This study has emphasized the occurrence of four communities colonizing gypsum outcrops belonging to the class *Psoretea decipientis* and to the orders *Barbuletalia unguiculatae* and *Tortulo brevissimae-Aloinetalia bifrontis*. The former order finds its optimum in the Mediterranean area up to Central European, the latter includes communities of arid and semi-arid areas of the Mediterranean and Irano-Turanian regions.

Tortuletum revolventis, *Trichostomo crispuli-Tortuletum revolventis* and *Crossidio squamiferi-Aloinetum aloidis* are included in the alliance *Grimaldion fragrantis*. In particular, *Tortuletum revolventis* is the most widespread community on the gypsum outcrops of Sicily, occurring in many sites in north-western, western and central Sicily. This pioneer association is known for the gypsum steppes of Thuringia (central Germany). *Trichostomo crispuli-Tortuletum revolventis* was found in north-western and central Sicily, showing more sciophytic, more mesic and more terricolous exigencies than *Tortuletum revolventis*. Both communities are exclusively found on gypsum substrate. *Crossidio squamiferi-Aloinetum aloidis* is a chasmophytic community, not strictly related to gypsum outcrops, having been found also on marl and clayey substrates.

Crossidio crassinervis-Tortuletum obtusatae, the only association referred to the order *Tortulo brevissimae-Aloinetalia bifrontis*, is the most xerophytic community among those found on the Sicilian gypsum outcrops; it is found only on gypsum crusts in conditions of high brightness and very high edaphic dryness, occupying favourable topographical positions for its survival. It represents a new record for the bryophyte vegetation of Italy.

The surveyed communities are rare and interesting from a naturalistic point of view. The peculiarity of the chalky substrate and the climatic conditions impose a strong selection on the species so that only these small and particular communities, characterized by species with peculiar xero-morphic adaptations, can survive. The almost exclusive occurrence of the life syndrome colonist with moderate rep hat the life syndromes and adaptive traits indicate the response of functional types towards environmental demands (e.g., Kürschner 2004, Kürschner and Frey 2013, Puglisi et al. 2016a, Puglisi et al 2016b).

Table 6. Life forms, life strategies and reproductive strategy of taxa (MPC values in %; abbreviations see Table 5).

Life forms	*Tortuletum revolventis*	*Trichostomo crispuli-Tortuletum revolventis*	*Crossidio squamiferi-Aloinetum aloidis*	*Cossidio crassinervis-Tortuletum obtusatae*
Tf	92.9	93.5	82.3	72.8
Ts	1.1	-	6.2	19.5
Tp	-	0.4	-	-
Tuft	4.9	4.2	11.6	-
We	0.1	1.5	-	1.2
Sc	1.0	-	-	6.5
St	-	0.4	-	-
Life strategy				
Colonist	90.7	94.7	95.7	91.0
Cas	11.9	5.6	-	8.7
Cs,as	3.6	-	9.2	-
Cs	13.0	5.9	28.1	7.2
Cm	62.2	83.2	58.4	75.1
Annual shuttle	1.4	-	-	7.8
Short-lived shuttle	0.2	0.4	-	-
Perennial shuttle	7.7	3.1	2.3	-
PeSs	7.1	0.1	0.3	-
PeSm	0.6	3.0	2.0	-
Perennial stayers	-	-	2.0	1.2
PSm	-	-	-	
PSas	-	-	-	
PSs	-	-	-	
PSs,as	-	-	2.0	

Many gypsum outcrops are included in protected areas, which are an important factor to guarantee the persistence of species and communities with legal protection. Despite this, it is assumed a reduction or even a high extinction probability of biodiversity, due to the effects of global warming with foreseen desertification phenomena in mid-latitude areas. Therefore, further conservation measures need to be implemented and the present study can give a contribution to the conservation and management of gypsum habitats, considered to be of high naturalistic value. For this purpose, it is important that the gypsum habitats of Sicily can be recognized as Habitats of European Community interest and referred to the Habitat 1520*, including the gypsophytic bryophyte associations Tortuletum revolventis, *Trichostomo crispuli-Tortuletum revolventis* and *Crossidio crassinervis-Tortuletum obtusatae*.

Syntaxonomic scheme

PSORETEA DECIPIENTIS Mattick ex Follmann 1974
BARBULETALIA UNGUICULATAE von Hübschmann 1960
Grimaldion fragrantis Šmarda & Hadàc 1944
Tortuletum revolventis Marstaller 1980
Community with *Trichostomum crispulum* and *Tortula revolvens*
Crossidio squamiferi-Aloinetum aloidis Guerra & Varo 1981
TORTULO BREVISSIMAE-ALOINETALIA BIFRONTIS (Ros & Guerra 1987) ex Puglisi 2010
Aloino bifrontis-Crossidion crassinervis Ros & Guerra 1987 ex Marstaller 2006
Crossidio crassinervis-Tortuletum obtusatae Ros & Guerra 1987

Acknowledgements

This research was financially supported by the research programme (PIA.CE.RI. 2020-2022 Line 2, cod. 22722132149) funded by the University of Catania.

Bibliography

Aleffi M, Pellis G, Puglisi M (2014) The bryophyte flora of six gypsum outcrops in the Northern Apennines (Nature 2000 Network, Emilia Romagna Region, Italy). Plant Biosystems 148(4): 825–836. https://doi.org/10.1080/11263504.2014.949331

Braun-Blanquet J (1964) Pflanzensoziologie - Grundzüge der Vegetationskunde, 3. Aufl. Springer, Berlin. 866 pp. https://doi.org/10.1007/978-3-7091-8110-2

Catalano R (1986) Le evaporiti Messiniane. Loro ruolo nell'evoluzione geologica della Sicilia. Le Grotte d'Italia 4(13): 109–122.

De Waele J, Piccini L, Columbu A, Madonia G, Vattano M, Calligaris C, D'angeli I. et al. (2017) Evaporite karst in Italy: a review. International Journal of Speleology 46(2): 137–168. https://doi.org/10.5038/1827-806X.46.2.2107

During HJ (1979) Life strategies of bryophytes: a preliminary review. Lindbergia 5: 2–18.

Escudero A, Palacio S, Maestre FT, Luzuriaga AL (2015) Plant life on gypsum: a review of its multiple facets. Biological Reviews of the Cambridge Philosophical Society 90(1): 1–18. https://doi.org/10.1111/brv.12092

Forti P, Sauro U (1996) The gypsum karst of Italy. International Journal of Speleology 25(3-4): 239–250. https://doi.org/10.5038/1827-806X.25.3.18

Frey W, Kürschner H (1991) Lebensstrategien von terrestrischen Bryophyten in der Judäischen Wüste. Botanica Acta 104: 172–182. https://doi.org/10.1111/j.1438-8677.1991.tb00214.x

Frey W, Herrnstadt I, Kürschner H (1990) Verbreitung und Soziologie terrestrischer Bryophytengesellschaften in der Judäischen Wüste. Phytocoenologia 19(2): 233–265. https://doi.org/10.1127/phyto/19/1990/233

García-Zamora R, Ros RM, Guerra (2000) Vegetación briofítica de las sierras de Filambres, Cabrera, Alhamilla y Cabo de Gata (Almería, SE de España). Cryptogamie Bryologie 21(1): 19–75. https://doi.org/10.1016/S1290-0796(00)00102-4

Guerra J, Ros RM, Martinez-Sanchez JJ, De Las Heras J, Herranz JM, Cano MJ (1993) Flora y vegetación briofítica de las zonas yesíferas de la provincia de Albacete. Al-Basit 32: 39-62.

Herrero J, Porta J (2000) The terminology and the concepts of gypsum-rich soils. Geoderma 96: 47–61. https://doi.org/10.1016/S0016-7061(00)00003-3

Hodgetts NG, Söderström L, Blockeel TL, Caspari S, Ignatov MS, Konstantinova NA, Lockhart N, Papp B, Schröck C, Sim-Sim M, et al. (2020) An annotated checklist of bryophytes of Europe, Macaronesia and Cyprus. Journal of Bryology 42(1): 1–116. https://doi.org/10.1080/03736687.2019.1694329

Kürschner H (2004) Life strategies and adaptations in bryophytes from the Near and Middle East. Turkish Journal of Botany 28: 73–84.

Kürschner H, Frey W (2013) Life strategies in bryophytes - a prime example for the evolution of functional types. Nova Hedwigia 96: 83–116. https://doi.org/10.1127/0029-5035/2012/0071

Lentini F, Carbone S (2014) Geology of Sicily. Memoria descrittive della Carta Geologica d'Italia - Volume XCV. pp. 7–414, figg. 533, tabb. 5; Tavv. 5.

Lo Giudice R, Bonanno G (2010) Bryophyte and Bryo-Tracheophyte diversity, life forms and life strategies in urban areas of Sicily. Nova Hedwigia 90(1–2): 161–194. https://doi.org/10.1127/0029-5035/2010/0090-0161

Mägdefrau K (1982) Life forms of Bryophytes. In: Smith A. J. E. (ed.), Bryophyte Ecology. Chapman & Hall, London. pp. 45–58. https://doi.org/10.1007/978-94-009-5891-3_2

Marstaller R (1980) Die Moosgesellschaften des Verbandes Phascion mitriformis Waldheim 1947. 7. Beitrag zur Moosvegetation Thüringens. Feddes Repertorium 91: 363–387. https://doi.org/10.1002/fedr.4910910512

Marstaller R (2008) Moosgesellschaften am Südrand des Kyffäusergebirges bei Bad Frankenhausen (Kyffäuserkreis). Mauritiana 20 (2): 289–348.

Moya JJ, Ros RM, Guerra J (1994) Bryophyte flora and vegetation of the Sierra del Maigmó Alicante, Spain. Cryptogamie, Bryologie-Lichénologie 15: 43–65.

Pesaresi S, Galdenzi D, Biondi E, Casavecchia S (2014) Bioclimate of Italy: application of the worldwide bioclimatic classification system. Journal of Maps 10: 538–553. https://doi.org/10.1080/17445647.2014.891472

Privitera M (1989) La vegetazione muscinale dei gessi dell'Agrigentino (Sicilia occidentale). Bollettino dell'Accademia Gioenia di Scienze Naturali Catania 22(335): 105–113.

Privitera M, Puglisi M (1999) Bryophyte vegetation of the badlands from the Aspromonte massif (S Italy). Nova Hedwigia 69(1–2): 195–210. https://doi.org/10.1127/nova.hedwigia/69/1999/195

Puglisi M, Privitera M (2012) A synopsis of the Italian bryophyte vegetation. Cryptogamie Bryologie 33(4): 357–382. https://doi.org/10.7872/cryb.v33.iss4.2012.357

Puglisi M, Kürschner H, Privitera M (2013) Saxicolous bryophyte communities of mountain areas of Greece: phytosociology, ecology, life forms and life strategies. Nova Hedwigia 97(1-2): 159–178. https://doi.org/10.1127/0029-5035/2013/0111

Puglisi M, Kürschner H, Privitera M (2014) The bryophyte association *Solorino saccatae-Distichietum capillacei* Reimers 1940 on the Mediterranean mountains: A tool for evaluating natural and conservative environments. Plant Biosystems 148(4): 815–824. https://doi.org/10.1080/11263504.2013.877092

Puglisi M, Kürschner H, Privitera M (2015) The *Schistidio atrofusci-Grimmietum pulvinatae*, a new association of the class Grimmietea anodontis on the Mediterranean mountains. Nova Hedwigia 100(1–2): 205–214. https://doi.org/10.1127/nova_hedwigia/2014/0213

Puglisi M, Kürschner H, Privitera M (2016a) The mountain and high-mountain bryophyte vegetation of the Pollino National Park (Southern Italy): syntaxonomy and ecology. Nova Hedwigia 103: 385–413. https://doi.org/10.1127/nova_hedwigia/2016/0362

Puglisi M, Minissale P, Sciandrello S, Privitera M. (2016b) Life syndrome of the bryophyte communities as an adaptative pattern in the Mediterranean temporary ponds of Italy. Pl Bios Plant Biosystems yst 150(6): 1417–1425. https://doi.org/10.1080/11263504.2016.1200688

Puglisi M, Kürschner H, Privitera M (2018) Phytosociology and life syndromes of bryophyte communities from sicilian caves, a clear example of relationship between bryophytes and environment. Plant Sociology 55(1): 3–20.

Puglisi M, Privitera M, Spampinato G (in press) The bryophyte vegetation of the gypsum outcrops: a focus on the communities with *Tortula revolvens*. Plant Biosystems.

Puglisi M, Campisi P, Privitera M, Spampinato G (2019) Bryophyte diversity in the gypsum outcrops of Sicily (Italy). Nova Hedwigia 111(3–4): 337–355. https://doi.org/10.1127/nova_hedwigia/2020/0602

Rivas-Martínez S, Rivas Sáenz S, Penas A (2011) Worldwide Bioclimatic Classification System. Global Geobotany 1: 1–634+4 Maps.

Ros RM, Guerra J (1987) Vegetación briofítica terrícola de la región de Murcia (Sureste de España). Phytocoenologia 15: 505–567. https://doi.org/10.1127/phyto/15/1987/505

Appendixes

Appendix I: Localities and dates of the relevés

Tab. 1 - Rel. 1: Serre di Ciminna (4 March 2019, 37°52'23.38"N, 13°35'28.21"E); rels 2, 3: Serre di Ciminna (4 March 2019, 37°52'24.2"N, 13°35'31.16"E); rels 4, 5: S. Ninfa (3 March 2019, 37°46'49.97"N, 12°53'50.65"E); rels 6, 7: S. Ninfa (3 March 2019, 37°46'49.58"N, 12°53'49.21"E); rel. 8: S. Ninfa (3 March 2019, 37°46'48.37"N, 12°53'54.00"E); rels 9-13: Sutera outskirts (from Privitera 1989); rels 14-19: Milena hills (from Privitera 1989); rels 20-25: Enna outskirts (from Lo Giudice and Bonanno 2010); rels 26-29: Raddusa outskirts (from Lo Giudice and Bonanno 2010); rels 30, 31: Castelluccio (4 March 2019, 37°46'20.63"N, 13°07'11.93"E).

Tab. 2 - Rels 1, 2: Castelluccio (4 March 2019; 37°46'19.5"N, 13°07'17.2"E); rels 3, 4: Castelluccio (4 March 2019; 37°46'13.6" N, 13°07'27.6" E); rels 5, 6:

Castelluccio (4 March 2019; 37°46'17.1" N, 13°07'33.7" E); Rels 7, 8: Sutera outskirts (7 April 2020; 37°30'01.5" N, 13°45'29.6" E); rel. 9: Sutera outskirts (7 April 2020; 37°30'02.8" N, 13°45'27.9" E); rels 10, 11 Sutera outskirts (7 April 2020; 37°30'06.2" N, 13°45'20.6" E); rels 12, 13: Milena hills (7 April 2020; 37°29'49.2" N, 13°45'02.4" E); rel. 14: Milena hills (7 April 2020; 37°29'52.1" N, 13°44'56" E).

Tab. 3 - Rel. 1: Sutera outskirts (6 April 2021, 37°30'20.34"N, 13°44'34.21"E); rel. 2: Sutera outskirts (6 April 2021, 37°30'19.66"N, 13°44'31.39"E); rel. 3: Sutera outskirts (6 April 2021, 37°30'19.34"N, 13°44'38.68"E); rel. 4: Milena outskirts (6 April 2021, 37°27'24.43"N, 13°44'37.66"E); rel. 5: Milena hills (6 April 2021, 37°27'23.90"N, 13°44'35.57"E); rel. 6,7: Raddusa outskirts (from Lo Giudice and Bonanno 2010).

Tab. 4 - Rels 1, 2: Rocca di Entella (4 March 2019; 37°46'20.72"N, 13°07'19.91"E); rel. 3: Rocca di Entella (4 March 2019; 37°46'20.20"N, 13°07'21.46"E); rel. 4,5: Rocca di Entella (4 March 2019; 37°46'21.37"N, 13°07'17.20"E); rels 6, 7: Rocca di Entella (4 March 2019; 37°46'17.96"N, 13°07'22.79"E); rel. 8: Rocca di Entella (6 April 2021; 37°46'19.77"N, 13°07'20.03"E).

Appendix II: Sporadic species

Table 1 – Rel. 1: *Cephaloziella baumgartneri* (1); rel. 2: *Aloina rigida* (+), rel. 4: *Weissia condensa* (1), *Entosthodon convexus* (+); rel. 5: *Tortella flavovirens* (1), *Aloina aloides* (+); rel. 6: *Aloina aloides* (1), *Weissia condensa* (1); rel. 11: *Bryum argenteum* (1); rel. 13: *Cheilothela chloropus* (1); rel. 14: *Ceratodon purpureus* (+), *Tortella flavovirens* (+); rel. 17: *Homalothecium sericeum* (+); rel. 18: *Timmiella barbuloides* (1); rel. 19: *Timmiella barbuloides* (1); *Homalothecium sericeum* (1); *Scleropodium touretii* (+); rel. 21: *Grimmia pulvinata* (+); rel. 23: *Bryum argenteum* (+), *Homalothecium aureum* (+), *Ptychostomum torquescens* (+); rel. 24: *Homalothecium aureum* (+), *Ptychostomum torquescens* (+); rel. 30: *Petalophyllum ralfsii* (+); *Lathagrium cristatum* (1); rel. 31: *Squamarina lentigera* (1).